## TUBERCULOSIS IN ARIZONA A Summary of Cases Reported through December 31, 1998

# Tuberculosis Control Section Office of Infectious Disease Services/Bureau of Epidemiology and Disease Control Arizona Department of Health Services

### **Morbidity**

Table 1: Tuberculosis Cases and Rates per 100,000 Population Arizona and United States, 1990-1998

		Arizona	United States			
Year Reported <sup>a</sup>	Cases	Population	Rate per 100,000	Cases d	Rate per 100,000 d	
1990	275	3,665,228 b	7.5	25,701	10.3	
1991	324	3,763,372 °	8.6	26,283	10.4	
1992	258	3,858,825 °	6.7	26,673	10.5	
1993	228	3,946,975 °	5.8	25,287	9.8	
1994	247	4,036,875 °	6.1	24,361	9.4	
1995	317	4,184,723 °	7.6	22,860	8.7	
1996	282	4,462,717 °	6.3	21,337	8.0	
1997	296	4,595,379 °	6.4	19,855	7.4	
1998	254	4,722,097 °	5.4	18,361	6.8	
Total	2,481	NA	NA	NA	NA	

- a Year reported reflects the year a case was counted by ADHS
- b Based on the 1990 Census
- c Based on population projections for the corresponding years
- d Data from the U.S. Department of Health and Human Services

#### **Demographics**

Table 2: Tuberculosis Cases and Rates per 100,000 Population by Age Group Arizona, 1998

100,000 Formation by Fige Group Hillzona, 1990									
Age Group	Cases # (%)	Population <sup>a</sup>	Rate per 100,000						
Under 5	11 (4.3)	351,612	3.1						
5 - 14	13 (5.1)	706,235	1.8						
15 - 24	21 (8.3)	647,480	3.2						
25 - 44	69 (27.2)	1,422,033	4.9						
45 - 64	72 (28.3)	934,149	7.7						
Over 64	68 (26.8)	660,588	10.3						
Total	254 (100)	4,722,097	5.4						

a Population data published by the ADHS Office of Epidemiology and Statistics (http://www.hs.state.az.us/plan/pop98/po98.htm)

Table 3: Tuberculosis Cases by Age Group and Sex Arizona, 1998

Age Group	Male	Female	Total Cases
Under 5	7	4	11
5 - 14	5	8	13
15 - 24	12	9	21
25 - 44	43	26	69
45 - 64	51	21	72
Over 64	42	26	68
Total	160	94	254

Table 4: Tuberculosis Cases and Rates per 100,000 Population by Age Group and Race/Ethnicity Arizona, 1998

Age Group	White, Not Hispanic	African-American	Hispanic, White	Native American	Asian/Pacific Islander
Under 5	2	1	7	1	0
5 - 14	0	1	12	0	0
15 - 24	2	3	12	2	2
25 - 44	16	8	32	7	6
45 - 64	27	2	22	16	5
Over 64	31	2	19	14	2
Total	78	17	104	40	15
1997 Population <sup>a</sup>	3,072,763	158,423	947,077	251,306	91,158
Rate per 100,000	2.5	10.7	11.0	15.9	16.5

a Race-specific rates based on 1997 data as 1998 data by race/ethnicity were unavailable as of April 16, 1999 (http://www.census.gov/population/www/projections/stproj.htm)

#### **Selected Clinical Characteristics**

Table 5: Tuberculosis Cases by Form of Disease and Vital Status at Diagnosis Arizona, 1998

Form of Disease	Cases Alive at Diagnosis	Cases Diagnosed after Death <sup>a</sup>	Total Cases
Pulmo nary Onl y b	186	6	192
Pulmo nary & Extra-pul monary b	17	0	17
Extra-pulmonary Only	41	4	45
Total	244	10	254

a It is unknown if TB was listed as a cause of death

Table 6: Tuberculosis Cases Resistant to INH and Other Anti-TB Drugs Arizona, 1993 - 1998

Year Reported	Cases	Culture Confirmed	Drug Sens Testing #	sitivity (%)	INH Resi	istant <sup>a</sup> (%) <sup>b</sup>	ME #	OR ° (%) b		Drug tance d (%) b	Total Drug I #	Resistance (%) b
1993	228	219	193	(88%)	11	(6%)	3	( 2%)	18	(9%)	32	(17%)
1994	247	223	204	(91%)	15	(7%)	4	( 2%)	17	(8%)	36	(18%)
1995	317	237	231	(97%)	8	(3%)	3	(1%)	29	(13%)	40	(17%)
1996	282	232	223	(96%)	6	(3%)	7	( 3%)	23	(10%)	36	(16%)
1997	296	235	223	(95%)	18	(8%)	2	(1%)	18	(8%)	38	(17%)
1998	254	208	204	(98%)	10	(5%)	1	(<1%)	14	(7%)	25	(12%)
Total	1,624	1,354	1,278	(94%)	68	(5%)	20	(2%)	119	(9%)	207	(16%)

a Resistant to at least INH without resistance to RIF

b The classification of Pulmonary TB includes laryngea l TB

b Percent of total cases with sensitivity testing

c Resistant to at least INH and RIF

d Resistance to drugs other than INH

#### **Risk Factor Information**

Table 7: Tuberculosis Cases by Selected Characteristics Arizona, 1993 - 1998

Risk Factor	1993 (%) a	1994 (%) a	1995 (%) a	1996 (%) a	1997 (%) <sup>a</sup>	1998 (%) <sup>a</sup>
<b>Total Cases Reported</b>	228	247	317	282	296	254
Occupation						
Correctional Facility Worker	0	0	0	0	0	1 (<1%)
Health Care Worker	3 (1%)	6 (2%)	8 (3%)	2 (1%)	3 (1%)	3 (1%)
Migrant Farm Worker	5 (2%)	7 (3%)	4 (1%)	9 (3%)	2 (1%)	4 (2%)
Reported Behaviors						
Injectable Drug User b	8 (4%)	4 (2%)	17 (5%)	12 (4%)	11 (4%)	6 (2%)
Non-injectable Drugs <sup>b</sup>	9 (4%)	5 (2%)	14 (4%)	13 (5%)	17 (6%)	10 (4%)
Excessive Alcohol Use <sup>b</sup>	22 (10%)	28 (11%)	43 (14%)	37 (13%)	45 (15%)	40 (16%)
Type of Residence						
Long Term Care Facility	5 (2%)	10 (4%)	16 (5%)	14 (5%)	12 (4%)	5 (2%)
Correctional Facility	6 (3%)	6 (2%)	16 (5%)	18 (6%)	23 (8%)	13 (5%)
Homeless <sup>b</sup>	31 (14%)	26 (11%)	45 (14%)	35 (12%)	27 (9%)	30 (12%)
Country of Birth						
Foreign Born	75 (33%)	88 (36%)	124 (39%)	108 (38%)	109 (37%)	100 (39%)
Underlying Disease						
Known HIV Infection <sup>c</sup>	16 (7%)	14 (6%)	28 (9%)	16 (6%)	12 (4%)	13 (5%)

a Percent of annual cases

Table 8: Tuberculosis Cases Among Foreign-born Persons by Length of Time in the United States Arizona, 1998

Area of Origin	Years in the United States Before Tuberculosis Report								
	<< 2< 2 Yea	rs2 - 32 - 3 Yea	ırs2 43YVanars	Unknown	Total Cases (%) <sup>a</sup>				
Asia/Pacific Islands	2	1	10	0	13 (13%)				
Mexico	16	7	45	1	69 (69%)				
Other Central/ South America & Caribbean	2	0	4	1	7 (7%)				
Europe and Former Soviet States	2	0	1	0	3 (3%)				
Africa	2	0	2	0	4 (4%)				
India	1	0	2	0	3 (3%)				
Unknown	0	0	1	0	1 (1%)				
Total	25	8	65	2	100 (100%)				

a Percent of total cases

b Within the year before TB diagnosis (routine collection of information began Jan. 1, 1993)

c HIV infection is a reportable condition in Arizona.

Table 9: Tuberculosis Cases by County of Residence and Other Selected Characteristics Arizona, 1998

County of Residence	Apache	Cochise	Coconino	Gila	Graham	Greenle e	La Paz	Maricopa	Mohave	Navajo	Pima	Pinal	Santa Cruz	Yavapai	Yuma	Total Cases
Cases	7	2	4	4	1	1	1	138	6	10	51	7	5	2	15	254
Rate/100,000 Population	10.6	1.7	3.4	8.5	3.0	11.3	5.2	4.9	4.4	11.5	6.2	4.6	13.6	1.4	11.4	5.4
Race/Ethnicity																
White, Not Hispanic	0	2	1	1	0	1	0	40	6	0	22	1	0	1	3	78
Hispanic, White	0	0	0	0	0	0	1	68	0	0	16	4	5	1	9	104
African-American	0	0	0	0	0	0	0	13	0	0	4	0	0	0	0	17
Native American	7	0	3	3	1	0	0	4	0	9	9	2	0	0	2	40
Asian/Pacifi c Islander	0	0	0	0	0	0	0	13	0	1	0	0	0	0	1	15
Sex/Age																
All Sexes <15 years	0	0	0	0	0	0	0	18	1	0	0	0	1	0	4	24
Females, 15 years	1	1	3	2	1	0	0	42	1	4	19	1	2	1	4	82
Males, 15 years	6	1	1	2	0	1	1	78	4	6	32	6	2	1	7	148
Type of Residence																
Correcti onal Facility	0	0	0	0	0	0	1	7	0	0	1	4	0	0	0	13
Long Term Care Facility	0	0	0	0	0	0	0	3	0	0	1	1	0	0	0	5
Homeless <sup>b</sup>	0	0	1	0	0	0	0	18	1	1	7	1	0	0	1	30
Country of Birth																
Foreign-born	0	0	0	0	0	0	1	66	1	1	15	4	4	1	7	100
Drug Resistance																
INH Resistance °	0	0	0	0	0	0	0	6	0	0	4	0	0	0	0	10
MDR <sup>d</sup>	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1

a Based on population data published by the Office of Epidemiology and Statistics (http://www.hs.state.az.us/plan/pop98/po98.htm)

File: 98stats.wpd/6May1999/dgm/seg

Database: 1998.dbf

b Within the year before TB diagnosis (routine collection of information began Jan. 1, 1993).

c Resistance to at least INH without resistance to RIF

d Resistance to at least INH and RIF